

GAS AND STARS IN AN HI-SELECTED GALAXY SAMPLE

JESSICA L. ROSENBERG^{1, 2}

Center for Astrophysics & Space Astronomy, Department of Astrophysical and Planetary Sciences, University of Colorado, Boulder, CO 80309

STEPHEN E. SCHNEIDER

Department of Astronomy, University of Massachusetts, Amherst, MA 01003

JENNIFER POSSON-BROWN²

Smith College, Northampton, MA 01063

Draft version February 2, 2008

ABSTRACT

We present the results of a J-band study of the HI-selected Arecibo Dual-Beam Survey and Arecibo Slice Survey galaxy samples using the 2 Micron All-Sky Survey data. We find that these galaxies span a wide range of stellar and gas properties. However, despite the diversity within the samples, we find a very tight correlation between luminosity and size in the J-band, similar to that we previously found (Rosenberg & Schneider 2003) between the HI mass and size. We also find that the correlation between the baryonic mass and the J-band diameter is even tighter than between the baryonic mass and the rotational velocity.

Subject headings: galaxies:fundamental parameters — galaxies: stellar content — infrared: galaxies

1. INTRODUCTION

In recent years several large “blind” 21 cm surveys for galaxies have been conducted providing a measurement of the gas in galaxies in the local universe (Meyer et al. 2004; Rosenberg & Schneider 2000; Zwaan et al. 1997; Spitzak & Schneider 1998). These surveys have revealed many galaxies with unusual characteristics: some without a definite optical counterpart and others with very high HI to stellar mass ratios, for example. These galaxy samples give us an opportunity to explore how the stellar properties of gas-rich galaxies differ from those of optically luminous systems.

Our understanding of the relationship between the global properties of gas and stars in galaxies has mostly been driven by studies of optically-selected, high surface brightness galaxies (e.g. Scodreggio & Gavazzi 1993; Huchtmeier & Richter 1985; Fisher & Tully 1981). Various efforts have been made to extend these studies to lower surface brightness galaxies, (e.g., Galaz et al. 2002, McGaugh et al. 2000, O’Neil et al. 2000, Sprayberry et al. 1995) revealing a great diversity of properties outside of the traditional “norms” defined by the high surface brightness samples, but such surveys remain tied to the requirement that the galaxies have formed stars in sufficient numbers and surface densities to be detected optically. Extragalactic HI surveys provide one of the few ways to probe the galaxy population independent of their luminosity and surface brightness. HI galaxy selection is also complementary to optical galaxy selection with respect to star formation history since the conversion of gas to stars renders a galaxy more visible optically, but less visible in an HI survey.

The blind HI surveys conducted with the Arecibo radio telescope remain among the deepest to date sampling galaxies with HI fluxes almost an order of magnitude smaller than the very large, but shallow HI Parkes All-Sky Survey (HIPASS, Meyer et al. 2004) and probing the lowest mass galaxies over a wider range of environments. The stellar properties of the Arecibo samples of Zwaan et al (1997) and Spitzak & Schneider (1998) have been studied at optical wavelengths. These studies have provided a picture of the gas-rich galaxies in the universe as containing a sub-sample of low-luminosity and low-surface-brightness galaxies (Spitzak & Schneider 1998) in higher proportion than found in optical surveys.

Infrared observations more nearly reflect the total stellar mass of galaxies, since they are less sensitive to the star formation rate and history and are less affected by dust than optical observations. In this paper we use the data from the 2 Micron All-Sky Survey (2MASS) to study the infrared properties of galaxies from the Arecibo Dual-Beam Survey (Rosenberg & Schneider 2000; RS) and from the Slice Survey (Spitzak & Schneider 1998; SS). Both of these surveys are “blind” HI surveys that probed the gas-rich galaxy populations in the local universe. The 2MASS observations of this galaxy sample are useful for examining the relationship between gas and stars in gas-rich galaxies in the local universe. While 2MASS provides information about the stellar mass in these galaxies, it suffers from a lack of surface brightness sensitivity which limits the galaxy detection rate. Nevertheless, over 85% of the galaxies in each sample were detected.

We discuss the HI and 2MASS data used in this study in §2 along with details about how some of the galaxies were measured from the 2MASS images. In §3 we discuss the relationship between the gaseous and stellar properties of the galaxies and discuss the surface densities of

¹ National Science Foundation Astronomy and Astrophysics Postdoctoral Fellow

² Present address: Center for Astrophysics, Cambridge, MA 02138

gas and stars in §4. In §5 we discuss the 2MASS images of the galaxies and in §6 we summarize our results.

2. DATA

2.1. Sample Selection - The Arecibo Dual-Beam and Slice Surveys

We examine the stellar properties of two HI-selected galaxy samples. Both the RS and SS surveys are “blind” 21 cm surveys carried out with the Arecibo 305 m telescope prior to the Gregorian upgrade. We have used these surveys to identify galaxies purely based on their gas content out to 7977 km s^{-1} and 8340 km s^{-1} respectively. The RS survey covered $\sim 430 \text{ deg}^2$ in the main beam and detected 265 galaxies while the SS survey covered $\sim 55 \text{ deg}^2$ and detected 75 galaxies. The selection functions and HI mass functions have been studied in detail and are presented in Rosenberg & Schneider (2002) and Schneider, Spitzak, & Rosenberg (1999) for the RS and SS surveys respectively.

The details of the calculations of line width, velocity, distance, and HI mass for the entire sample are presented in RS. The SS survey additionally includes broadband B, R and I optical data. There are three galaxies from the RS survey for which the RS line widths differed from the literature values because one horn of the velocity profile was missed in the survey measurement. In these cases (RS 17, RS 109, and RS 189) we use the values from the Huchtmeier & Richter catalog (1985) when using line widths to calculate the dynamical mass (see §3).

2.2. The 2-Micron All Sky Survey Data

The 2 Micron All-Sky Survey (2MASS) provides simultaneous J, H, and K_s -band observations of the entire sky; the 2MASS project processed the data to generate a point source catalog and an extended source catalog. We use the extended source catalog data for the RS and SS galaxies in the following analyses and also supplement this with our own analysis of the images for a number of galaxies that were undetected by the standard processing procedures (discussed in §2.2.1). A full description of the galaxy detection algorithm and the resulting catalogs are available in the Explanatory Supplement to the 2MASS All Sky Data Release³ (Cutri et al.).

The 2MASS extended source catalog has undergone several iterations and we use results from both the Version 2 and Version 3 catalogs in these analyses. We use Version 2 in addition to Version 3 because this earlier version of the software used previous galaxy catalog positions to seed the search algorithms (in addition to doing an independent automated search) and therefore measured a number of fainter sources that Version 3 did not detect, since it used only the automated algorithms. In addition, there are some large galaxies for which we report the Version 2 values rather than the Version 3 values because some of the information is missing in the version 3 catalog.

³ <http://www.ipac.caltech.edu/2mass/releases/allsky/doc/explsups.html>
by Cutri, R.M., Skrutskie, M.F., Van Dyk, S., Beichman, C.A., Carpenter, J.M., Chester, T., Cambresy, L., Evans, T., Fowler, J., Gizis, J., Howard, E., Huchra, J., Jarrett, T., Kopan, E.L., Kirkpatrick, J.D., Light, R.M., Marsh, K.A., McCallon, H., Schneider, S., Stiening, R., Sykes, M., Weinberg, M., Wheaton, W.A., Wheelock, S., Zacharias, N.

The J-band observations are the most sensitive of the three 2MASS bands and our HI-selected sources tend to be blue on average, so we use these data in our analyses. The catalog data were used for 45 (4 from Version 2, 41 from Version 3) of the 75 SS galaxies and 180 (50 from Version 2, 130 from Version 3) of the 265 RS galaxies. Some of the galaxies that were not in the extended source catalogs can be identified on the full resolution 2MASS images; in these cases we measured the galaxy from the image. Details of our image analysis are described below. There were an additional 19 SS and 47 RS galaxies that we measured from the images. For 11 of the SS galaxies and 38 of the RS galaxies there was no cataloged detection and we were not able to measure the galaxy from the images. For most of the galaxies that were not measured, the source was just too faint to be detected in the short 2MASS exposures, but some exceptions are described in the specific galaxy notes below.

The near infrared measurements for the galaxies in both the RS and SS surveys are given in Tables 1 and 2 in the Appendix. The tables also list the source for the measurements, i.e., whether the data come from Version 2 or 3 of the 2MASS catalog, were measured from the image, or if the source was not detected.

2.2.1. 2MASS Measured Data

To measure the galaxies on the 2MASS images we have used the ELLIPSE package in IRAF. The brighter galaxies were interactively fit with ellipses such that the ellipse parameters were allowed to change in the highest surface brightness regions but when the fits became uncertain in the outer regions, the ellipse parameters were held fixed while additional steps in radius were taken. For the lowest surface brightness galaxies, even the central regions had uncertain parameter fits, so only fixed circular apertures were used for the photometry.

ELLIPSE was run at least twice for each galaxy so that the outer “background” ellipses could be used for background subtraction. Then the fitting was rerun on the background-subtracted image. The background subtraction was repeated if an adequately flat background had not been obtained on the first pass. In all cases, the point sources in the image were masked prior to the ellipse fitting.

2.2.2. Notes About Selected Galaxies

We provide information about the near infrared measurements for a few galaxies that warrant a comment. The numbers refer to the entry number in the RS survey catalog (see Table 1 for the ADBS names).

- **RS 50:** The HI detection of this system includes an interacting pair of galaxies. Since we do not separate the pair in the HI observation we do not try to correlate the HI with a near infrared measurement of one galaxy or the other.
- **RS 73:** We do not report an infrared measurement for this galaxy because it is too contaminated by a bright star.
- **RS 79:** We use the Version 2 measurement of the brighter galaxy in the interacting pair. Both galaxies are probably contributing to the HI, but includ-

ing the fainter of the two galaxies would change the J-band magnitude by less than 0.3 mag.

- **RS 184:** This galaxy is in the Version 3 extended source catalog. However, we use measurements from the image because the center of this very low surface brightness galaxy was missed in favor of a field star by the galaxy detection algorithm.

2.3. Using the 2MASS data as a Measure of Stellar Mass

As a basis for understanding the properties of these HI-selected galaxies we want to compare the gas mass and the stellar mass in these systems. The 2MASS data are particularly good for measuring the stellar mass as they are more sensitive to low mass stars and less sensitive to dust and star formation than optical observations. However, there is substantial scatter in the relationship between luminosity and stellar mass even in the infrared. The scatter can be decreased by using information about the galaxy's color (Bell & de Jong 2001).

Based on stellar population synthesis models, a variety of estimates have been made for various stellar initial mass functions and star-formation histories. Bell & de Jong (2001) present a range of models that yield realistic colors and surface brightnesses for present-day galaxies. Bell (private communication) has provided us with J-band values for these same models and finds a tight correlation between (B-R) color and the ratio of stellar mass to J-band luminosity:

$$\log(M_{star}/L_J) = 0.552(B - R) - 0.724 \quad (1)$$

The value of M_{star}/L_J varies from ~ 0.4 for blue galaxies $[(B - R) \sim 0.6]$ to ~ 1.2 for red galaxies $[(B - R) \sim 1.6]$. Including differences between models and measurement uncertainties, there is thus a range of almost a factor of 4 in the relationship between stellar mass and J-band luminosity for the normal range of galaxy colors. When the color is known, the total range reduces to about a factor of 1.5. These results appear to be consistent with alternative models carried out for I and H bands (cf. McGaugh & de Blok 1997; McGaugh et al. 2000).

The SS galaxies were previously observed in broadband colors, which allows us to estimate their mass-to-light ratios more precisely. Their average (B-R) color was 1.0, so their average M_{star}/L_J value is 0.7. Figure 1 shows the relationship between M_{star}/L_J for the SS galaxies calculated using Equation 1 and their ratio of HI mass to J-band luminosity, M_{HI}/L_J . The total neutral hydrogen mass was calculated from the published data using the well-known relationship $M_{HI} = 2.36 \times 10^5 D^2 I_{HI}$, where D is the distance in Mpc and I_{HI} is the integrated flux in Jy km s^{-1} . The standard deviation around the line is 0.15 while the standard deviation around the average value of $\log(M_{star}/L_J) = -0.15$ is 0.19. The correlation is not extremely tight, but it allows us to improve our estimate of the mass-to-light-ratios for the galaxies that do not have broadband colors: the RS galaxies and the faintest SS galaxies.

In addition to the intrinsic and measurement uncertainties in determining M_{star}/L_J , an additional source of scatter is introduced in the determination of stellar




FIG. 1.— The relationship between the mass-to-light ratio calculated using Equation 1 and the M_{HI}/L_J for the SS data. The fit to these data are used to improve the estimate of M_{star}/L_J for the data for which B-R measurements are not available.

mass because the 2MASS data are not very deep. Uncertainties in the J-band flux are caused by the difficulty in measuring the isophotal sizes near the limit of the survey surface brightness sensitivity, particularly for the lowest surface brightness galaxies. We compare the isophotal measurements of galaxy magnitudes at $21 \text{ mag arcsec}^{-2}$ to the measurements in the largest aperture before the noise takes over (i.e., before the values start oscillating between fainter and brighter values due to noise). For the 2MASS cataloged values, we use the brightest aperture value reported. To be consistent with the catalog measurements, we restrict our own measured apertures to those used in the catalog (i.e., 5, 7, 10, 15, 20, 25, 30, 40, 50, 60, or 70 arcsec). However, 19 of the RS galaxies are significantly larger than $70''$ so the largest aperture is not a good measurement of its size of flux. For these galaxies, we use the extrapolated J-band magnitude so that the size and flux are not severely underestimated. Table 1 indicates for which galaxies the extrapolated values are used. There are no SS galaxies that are significantly larger than the $70''$ aperture.

Figure 2 shows that the magnitudes for most of the faint galaxies are severely underestimated if the isophotal value is used. For all of the J-band luminosities in this paper we use the aperture measurement values. Additionally, we adopt a value of $M_{\odot}(J) = 3.73$ (Johnson 1966; Allen 1973) for our conversion to solar luminosities.

3. STARS AND GAS IN AN HI-SELECTED SAMPLE

By using the RS and SS data, we have selected a wide range of galaxy types with the only criteria being that they contain neutral hydrogen. These systems span the range from dwarfs and irregulars to early-type spirals and interacting systems. Appendix A shows the 2MASS near infrared images for these galaxies; their morphology is discussed in greater detail in §5.

Figure 3 shows the relationship between the J-band surface brightness and M_{HI}/L_J for the RS galaxies (filled circles) and for the SS galaxies (open circles). The J-




FIG. 2.— The relationship between aperture and 21 mag arcsec⁻² isophotal magnitudes from the RS and the SS samples. The open symbols indicate magnitudes that were obtained from the 2MASS extended source catalog, Version 2 or 3. The closed symbols indicate magnitudes that were measured from the images.




FIG. 3.— The relationship between the average J-band surface brightness and the ratio of HI mass to J-band luminosity for the RS galaxies (filled circles) and the SS galaxies (open circles).

band surface brightness is determined within the aperture radius (or extrapolated radius for the largest galaxies) determined as discussed in the previous section. The surface brightness that we derive is, roughly, a surface brightness at a slightly fainter isophote than $J=21$ mag arcsec⁻². This figure illustrates that, in general, the gas-rich galaxies (galaxies with high values of M_{HI}/L_J) are the lower surface brightness ones indicating that HI-selection is a good way to find low surface brightness galaxies. Despite not having a very precise measure of surface brightness, the correlation with M_{HI}/L_J is pretty good.

The typical (B-J) color for a spiral galaxy is between 2 and 3. Using (B-J) = 2 as a conservative value, the

average surface brightnesses of these HI selected galaxies range between $\Sigma_J = 18$ to 25 mag arcsec⁻². There are a variety of different definitions of what constitutes a low surface brightness galaxy. One definition refers only to the central surface brightness of the disk component after a bulge-disk decomposition has been carried out. By this definition, many galaxies with high surface brightness bulges are categorized as LSB, because the disk component is faint. Another common definition uses the mean blue surface brightness within the $\mu_{B_0} = 25.0$ mag arcsec⁻² isophote, giving the label of LSB to galaxies with inclination-corrected mean surface brightnesses dimmer than $\langle \mu_{B_0} \rangle > 25.0$ mag arcsec⁻².

Since we cannot perform bulge-disk decompositions with the present data, our mean J-band surface brightnesses within the $\mu_J = 21.0$ mag arcsec⁻² are most readily comparable to the latter definition, although for typical (B-J) colors we measure the mean within a brighter isophote. It appears, though, that analogous to the B-band definition of LSB, we might call galaxies dimmer than $\langle \mu_J \rangle > 21.0$ mag arcsec⁻² infrared LSBs. This cut-off for LSBs also marks the approximate point at which the total stellar luminosity becomes smaller than the HI mass (in solar units).

The total baryonic mass was estimated from the HI and J-band emission using:

$$M_{bar} = M_{star} + 1.4 \times M_{HI}. \quad (2)$$

where M_{star} is the total stellar mass, derived from the J-band luminosity mass-to-light ratio discussed in the previous section, and the total gas mass is estimated from M_{HI} multiplied by 1.4 to account for helium and metals. Obviously, including an estimate of the molecular and ionized gas would increase this further. The stellar and gas masses both have significant uncertainties associated with them (e.g., due to the metallicity of the stars and the amount of molecular gas present), but the effect of internal extinction on the estimate is small compared to its effect at optical wavelengths since the extinction at J is only about 20% of that at B. Furthermore, the additional gas mass is likely to be largest for earlier type galaxies (Young & Knezek 1991), which are dominated by their stellar content, and the uncertainties in the stellar mass are largest for the dim galaxies that are generally dominated by their neutral hydrogen mass, so the uncertainties should have little impact on the total baryonic mass estimates.

Figure 4 shows how the various measured and derived values of mass and luminosity vary with M_{HI}/L_J . We find that the HI mass is essentially uncorrelated with the ratio for our HI-selected sample; consequently, the stellar luminosity declines roughly in inverse proportion to the M_{HI}/L_J ratio. An indication of dynamical mass was calculated for our galaxies using their HI line widths (see RS and SS for a discussion of the measurements of line width) and optical dimensions to estimate the inclination and rotation speed, using the equation:

$$M_{dyn} = v_{rot}^2 r_{opt} / G. \quad (3)$$

v_{rot} is the inclination corrected line width ($v/\sin[i]$). We limited the inclination correction to inclinations greater than $i > 66^\circ$ ($b/a \leq 0.4$). This calculation of dynamical mass will not accurately estimate the masses




FIG. 4.— The dependence of stellar luminosity, HI mass, total baryonic mass, and dynamical mass upon the ratio of gas to stars. The distance independent quantities of the ratio of dynamical mass to J-band luminosity and the ratio of dynamical mass to baryonic mass are also plotted as a function of the gas to stars ratio. The filled circles are the RS galaxies, the open circles are the SS galaxies. The triangles plotted in panels containing dynamical mass are the galaxies for which $b/a > 0.4$ for which the dynamical mass measurements may not be reliable.

of dwarf systems that are not primarily rotationally supported, and this mass estimate is directly dependent on the limits to which starlight is seen, but will, nevertheless give an indication of the total mass inside the faintest detected isophote. We have chosen to use the optical size in this calculation, despite the fact that there is often significant HI mass outside of this radius, because HI sizes are only available for a small subset of the RS data and are not available for any of the SS data. The optical sizes of the RS galaxies are derived from Palomar Sky Survey images and represent roughly the B-band 25 mag arcsec⁻² isophotal size. In Figure 4 the galaxies with $b/a > 0.4$ (more face-on systems for which the determination of dynamical mass is much less reliable) have been plotted as triangles.

Even though there are some clear trends in the properties of the galaxies in these surveys, Figure 4 also illustrates an important diversity. The width of the distribution of masses, luminosities and mass-to-light ratios at any given value of M_{HI}/L_J is many times larger than might be caused by measurement uncertainties alone. The spread is probably even larger than these figures show since the lower-mass sources were only detectable to a limited distance within our search volume. What Figure 4 shows is that galaxies of any given HI mass span the entire range of gas-to-star ratios – we see galaxies that are low in M_{HI} because they have turned most of their gas into stars and some that have low M_{HI} because they are low mass systems with little of their mass in stars.

We note the scarcity of galaxies with $M_{HI}/L_J \lesssim 0.03$ in Figure 4. It is not clear whether this scarcity is a selection effect or a true deficit of galaxies. Presumably, some ellipticals and lenticulars would fall in this range, and some early-type systems *were* detected in these surveys. The total contribution of early type galaxies containing HI is




FIG. 5.— The relationship between the J-band luminosity and HI mass for the RS (filled circles) galaxies and the SS (open circles) galaxies. The dashed lines indicate M_{HI*} (Rosenberg & Schneider 2002) and L_{J*} (from the 2dF sample using 2MASS measurements; Cole et al. 2001). The solid line indicates a one-to-one relationship between the parameters.

uncertain. At such low M_{HI}/L_J values, the HI selection criteria only permit us to detect high stellar luminosity sources, and even those only out to a relatively small distances. HI data for an unbiased optical sample is needed to determine their true statistical contribution.

The large range of galaxy properties and M_{HI}/L_J values for the RS and SS surveys indicates that there is not an easy conversion between the HI mass function and the J-band luminosity function. Figure 5 further demonstrates the problem with using one of these parameters to predict the other. It is not clear that there is a linear relationship between L_J and M_{HI} at all masses and luminosities and the spread around the correlation between the quantities is large. As we noted in our earlier paper (Rosenberg & Schneider 2002), at a given HI mass, there may be several orders of magnitude variation in the J-band luminosity, making HI mass a poor predictor of stellar content, and vice versa.

Both the baryonic and dynamical mass estimates are higher for sources with smaller ratios of M_{HI}/L_J . As we show in Figure 4, however, the dynamical mass-to-light ratio is higher on average for the gas-rich galaxies. This figure also shows that the ratio of dynamical mass to baryonic mass is relatively flat as function of M_{HI}/L_J with a median value of 3.8 for galaxies with $b/a < 0.4$. The range in values that that we find for M_{dyn}/M_{bar} is comparable to the range for disk galaxies (Zavala et al. 2003). Even excluding the more face-on systems for which the dynamical mass calculation is problematic, there are 5 galaxies with dynamical masses that are smaller than their baryonic mass estimate. For 2 of the RS cases, the measurement of inclination is for the bright central region, but there is a more diffuse light distribution that may indicate that these systems are more face on. For the other 3 systems it is not clear why the dynamic mass is much smaller than the baryonic mass except that they seem to have low rotation velocities for such edge-on systems that might indicate that the line




FIG. 6.— The baryonic mass for the ADBS galaxies (filled circles) and the AS galaxies (open circles) plotted against the dynamic mass, rotational velocity, and J-band size. The triangles in the dynamic mass and rotational velocity plots indicate the galaxies for which $b/a \geq 0.4$ and are, therefore, less reliable. The solid lines in the plots show the average of forward and backward least squares fits to the data excluding the galaxies with $b/a \geq 0.4$.




FIG. 7.— The scatter of the baryonic masses around the least squares fits to each of the quantities shown in Figure 6. The rms dispersion of the fit is noted at the top of each panel.

width has not been properly measured (e.g., missing one horn of a double horned HI profile).

Figure 6 shows the relationship between our estimate of the baryonic mass and a variety of other measurements that relate to the overall mass of a galaxy: the dynamical mass (as described in §3), the rotation speed, and the radius. Note that we have marked the face-on ($b/a \geq 0.4$) galaxies with triangles and have not included them in the fits.

It is apparent by eye that the scatter is significantly smaller for galaxy radius than for other more traditional estimators of mass, although when we remove the more face-on galaxies from our calculation, the difference between the dynamical mass and J-band size is greatly reduced. We averaged forward and backward linear least squares fits to these data and display the distribution around the fit and the calculated sigma in Figure 7. Thus, the size of a galaxy makes a good predictor of the mass present. While galaxy models do indicate that size might be the third parameter in a spiral galaxy fundamental plane relationship (Shen, Mo, & Shu 2002), the physics behind this correlation is unclear.

4. THE SURFACE DENSITY OF MATTER IN GALAXIES

Despite the diverse range in galaxy properties discussed in the previous sections, there is a tight correlation between HI mass and HI size (Rosenberg & Schneider 2003) as well as a good correlation between J-band luminosity and J-band size. We examine the mass surface densities of gas, stars, and baryons since the Schmidt Law associates the surface density of gas in a galaxy with




FIG. 8.— The J-band luminosity for the RS galaxies (filled circles) and the SS galaxies (open circles) plotted against the dynamical mass, rotational velocity, and J-band size. The triangles in the dynamical mass and rotational velocity plots indicate the galaxies with $b/a \geq 0.4$. The solid lines show the forward and backward least squares fits to the data excluding galaxies with $b/a \geq 0.4$.

the star formation rate (Kennicutt 1998) and seems to imply a regulation mechanism between the gas and the star formation that might help explain the relationship between mass and size for these two populations.

Figure 8 shows the relationship between luminosity and dynamical mass, rotational velocity, and J-band size for these galaxies. Forward and backward linear least squares fits to these data are plotted. The standard deviation around these fits are 0.49, 0.69, and 0.37 respectively, providing an indication of the significance of the R_J versus L_J correlation. For the dynamical mass and rotational velocity correlations we have plotted the galaxies with axis ratios greater than 0.4 as circles (galaxies with axis ratios of 0.4 or less are plotted as triangles). We have only included those galaxies with axis ratios greater than 0.4 in these two fits.

The relationship between J-band luminosity and J-band size is tighter than the Tully-Fisher relationship (Tully & Fisher 1977) for this sample which includes irregulars, interacting systems, and dwarfs as well as spirals. Unfortunately, since luminosity and size scale the same way with distance, the tight correlation does not offer any possibilities for distance estimation.

The Tully-Fisher relation (Tully & Fisher 1977) has provided an important test of galaxy formation simulations and has been very hard to reproduce in detail; most of the galaxies produced in simulations lose too much of their angular momentum without large amounts of ad hoc energy injection into the system. Governato et al. (2004) have managed to produce one galaxy with the correct dynamical properties by going to higher resolution in their simulations, but still has an excess of massive satellites and a paucity of cold gas. Even as this work shows a success in producing the dynamical properties of a galaxy it highlights how much is left to be done. The Tully-Fisher relation seems to indicate that there is a physical connection between the dark matter component and the baryonic component in spiral galaxies.

For these HI-selected galaxies, many of which are dwarfs or irregulars rather than spirals, we find less of a correlation between V_{rot} or M_{dyn} and M_{bar} than is usually found for spiral galaxies (e.g., Giovanelli et al. 1997) culling out the systems with axis ratios greater than or equal to 0.4. However, even for this disparate group of galaxies, the average surface density of gas (Rosenberg & Schneider 2003) has a very small range relative to the variation in the stellar surface density. The surface density of the gas is determined as the HI mass divided by




FIG. 9.— The distribution of mass surface densities of gas, stars, and baryons. The gas surface densities, indicated by the dark-shaded histogram, come from the RS sample galaxies with measured HI sizes. The open histogram shows the stellar mass surface densities of the RS and SS galaxies using the J-band luminosity to estimate stellar mass, and the J-band sizes. The total baryonic mass surface density (cross-hatched histogram) is estimated using the sum of gas and stars and the optical size of each galaxy.

the HI area defined by the size at $2 \times 10^{20} \text{ cm}^{-2}$.

Figure 9 shows the HI surface densities for the RS galaxies with measured HI sizes. Rosenberg & Schneider (2003) showed that the average surface density of the gas in these galaxies was nearly constant, and this result is evident in the small range in mean HI surface densities: the values range from 3 to $24 \text{ M}_{\odot} \text{ pc}^{-2}$ with only 4 of the 50 galaxies having values above 10. We do not discuss the surface densities of the SS sample because HI sizes are not available for these galaxies. Alternatively, the stellar surface densities or the RS and SS sample cover a broad distribution ranging from 0.8 to $660 \text{ M}_{\odot} \text{ pc}^{-2}$. The notable feature of this plot is that the dispersion in the HI surface density is only 0.46 while the dispersion around the stellar mass surface density is 4.5. The baryonic mass surface density is the combination of the HI and stellar masses as given in Equation 2 and has a standard deviation around the mean surface density (averaged over the HI size) of 5.8. These mass surface densities may provide an additional constraint for the simulations.

The larger dispersion in the stellar mass surface density of the RS and SS samples is, at least in part, due to the large uncertainties in the J-band sizes. Also, since the diameters were measured at a high surface brightness, the mean surface densities are higher than they would have been had we measured out to an equivalently low surface density as the gas. The median value for the distribution is $19.3 \text{ M}_{\odot} \text{ pc}^{-2}$ relative to $5.9 \text{ M}_{\odot} \text{ pc}^{-2}$ for the HI mass density. We note that the median value of the mass surface density is generally higher for the stars than for the gas. This corresponds well with the observation that many gas-rich galaxies have gas distributions extending well beyond the optical (or near infrared) light distributions. However, we do note that there are still some systems in the sample that have very low stellar

mass surface densities providing an indication that we have detected some very low surface brightness systems.

5. THE MORPHOLOGY OF HI-SELECTED GALAXIES

Figures 1 and 3 in the Appendix show the 2MASS J-band images for the RS and SS galaxies with measured values of baryonic mass, respectively. The galaxies have been placed in order of their baryonic mass from the highest baryonic mass systems to the lowest (within each survey). Figures 2 and 4 show the galaxies for which we do not measure baryonic mass.

The images of these galaxies illustrate the diverse population as discussed in §3. The sample covers the range from nearby bright spiral galaxies like RS 189, NGC 4565, to bulge-less low surface brightness smudges like RS 184 (Figure 1 in the middle of the second to last line on the second page). There are also early type galaxies in the sample like SS 19 (Figure 3 in the middle of the last line) which is NGC 7712, an elliptical galaxy (Scodeggio et al. 1995) and close interacting pairs like RS 50 (second to last image in Figure 2).

The SS98 optical images of the SS sample indicate that a substantial fraction of the galaxies are low surface brightness. As discussed in §3, these 2MASS data also indicate that there is a substantial population of low surface brightness galaxies among the RS and SS samples even though the surface brightness can not be directly compared with the usual low surface brightness definition. Comparing the 2MASS images with the SS98 optical images, we find that nearly all of the low surface brightness galaxies are not detected or are barely detected in the 2MASS data. Inevitably this is because 2MASS is a shallow survey, but there is no evidence for bulges that are bright in the infrared but not in the optical as was found for low surface brightness galaxies by Galaz et al. (2002).

Visual inspection of the RS galaxy images seems to indicate that most of the highest baryonic mass galaxies are spirals with prominent bulge components while the lower baryonic mass galaxies show more variation galaxy type and bulge size down to galaxies like RS 184 where the brightness distribution appears fairly flat across the J-band detected stellar disk. This same trend is not apparent for the SS galaxies, but the sample size is significantly smaller which might account for the difference.

6. SUMMARY

We have examined the stellar properties of two HI-selected galaxy samples and found that they show a large range of stellar properties. We find that the galaxies cover a wide range in mass-to-luminosity ratio and the ratio is uncorrelated with the HI mass of the system. The range suggests that star formation does not proceed uniformly in all galaxies, some of the galaxies have funneled a large fraction of their gas mass into stars by the present day while there are other galaxies that are still largely dominated by their gas content. Because of these differences, one cannot infer the gas properties of a galaxy from its stellar properties and vice versa. The different proportions of gas and stars in these galaxies may be providing us a glimpse of galaxies in different stages of evolution.

Despite the wide range of gas-to-star ratios for galaxies in these samples, there are some surprising correlations.

There is a small range in the average HI surface density in these galaxies. While the average stellar surface density distribution is not as tight, it is also a fairly narrow distribution given all of the inaccuracies in its measurement. The baryonic matter appears to average to about one quarter of the dynamical mass (within the optical dimensions of the galaxy) across a wide range of galaxy types.

Most notably, the size of the stellar disk appears to be a very good predictor of the total baryonic content of galaxies. The physical causes of this correlation are not immediately apparent, but the correlation spans over three orders of magnitude in galaxy mass.

We would like to thank 2MASS for the effort that has gone into this catalog and for providing financial support for this research. We would particularly like to thank Roc

Cutri for all of the help in obtaining the full resolution images that were needed for this paper to be possible. We would also like to thank the Arecibo and VLA staffs for their assistance with the HI observations. Thanks also to Eric Bell for the information on J-band M/L ratios. We also appreciate John Spitzak's work on the program to display and print the galaxy images. Thanks also go to the anonymous referee for helpful suggestions. JLR acknowledges support from the National Science Foundation under grant AST-0302049. The Digitized Sky Surveys were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates were processed into the present compressed digital form with the permission of these institutions.

APPENDIX

In this appendix we present the 2MASS data for the HI-selected samples. The data include tables of the 2MASS size and magnitude values which are presented since many of these values are measured from the images themselves or are derived from the Version 2 extended source catalogs which are not generally available. The information in Table 1, the ADBS data, is as follows: (1) the sequential listing in the RS catalog; (2) the name from the ADBS catalog; (3, 4) the RA and DEC from the HI catalog, (5,6) the RA and DEC from the 2MASS catalog if it was identified or of the 2MASS image center, (7) the radius of the 21 mag arcsec⁻² isophote; (8) the b/a axis ratio used to measure the magnitude within the 21 mag arcsec⁻² isophote; (9) J 21 mag arcsec⁻² isophotal magnitude (J_{21}); (10) Error in J_{21} , (11) the radius used for the aperture magnitude; (12) J-band aperture magnitude; (13) error in the J-band aperture magnitude; (14) method used for determining the magnitudes – V2 and V3 refer to sources measured from the Version 2 or 3 catalog, MI are sources measured off the images, and NM are sources for which there was nothing on the 2MASS images to measure. The data in Table 2, the SS data, are the same except that the SS catalog name and the sequential catalog number are the same so all columns starting from the RA and DEC from the HI catalog are shifted one to the left.

The images are presented as a quick look at the galaxies in the sample. The objects have been placed in order of baryonic mass from the most massive to the least massive. Objects in Figures 2 and 4 are sources for which a baryonic mass was not determined.

REFERENCES

- Allen C. W., 1973, *Astrophysical Quantities*, 3rd edn. Athlone Press, London.
- Bell, E. F. & de Jong, R. S. 2001, *ApJ*, 550, 212.
- Cole, S., Norberg, P., Baugh, C. M., Frenk, C. S., Bland-Hawthorn, J., Bridges, T., Cannon, R., Colless, M., Collins, C., Couch, W., Cross, N., Dalton, G., De Propris, R., Driver, S. P., Efstathiou, G., Ellis, R. S., Glazebrook, K., Jackson, C., Lahav, O., Lewis, I., Lumsden, S., Maddox, S., Madgwick, D., Peacock, J. A., Peterson, B. A., Sutherland, W., Taylor, K. 2001, *MNRAS*, 326, 255.
- Fisher, J. R. & Tully, R. B. 1981, *ApJ*, 243, L23.
- Galaz, G., Dalcanton, J. J., Infante, L. & Triester, E. 2002, *ApJ*, 124, 1360.
- Governato, F., Mayer, L., Wadsley, J., Gardner, J. P., Willman, B., Hayashi, E., Quinn, T., Stadel, J., & Lake, G. 2004, *ApJ*, 607, 688.
- Huchtmeier, W. K. & Richter, O. -G. 1985, *A&AS*, 149, 118.
- Johnson H. L., 1966, *ARA&A*, 4, 193.
- Kennicutt, R. C. Jr. 1998, *ApJ*, 498, 541.
- McGaugh, S. S., Schombert J. M., Bothun, G. D., & de Blok, W. J. G. 2000, *ApJ*, 533, 102.
- McGaugh, S. S. & de Blok, W. J. G. 1997, *ApJ*, 481, 689.
- Meyer, M. J., Zwaan, M. A., Webster, R. L., Smith, L., Ryan-Weber, E., Drinkwater, M. J., Barnes, D. G., Howlett, M., Kilborn, V. A., Stevens, J., Waugh, M., Pierce, M. J., Bhathal, R., de Blok, W. J. G., Disney, M. J., Ekers, R. D., Freeman, K. C., Garcia, D. A., Gibson, B. K., Harnett, J., Henning, P. A., Jerjen, H., Kesteven, M. J., Knezek, P. M., Koribalski, B. S., Mader, S., Marquarding, M., Minchin, R. F., O'Brien, J., Oosterloo, T., Price, R. M., Putman, M. E., Ryder, S. D., Sadler, E. M., Stewart, I. M., Stootman, F., Wright, A. E. 2004, *MNRAS*, 350, 1195.
- Mo, H. J. & Mao, S. 2000, *MNRAS*, 318, 163.
- O'Neil, K., Bothun, G. D., & Schombert, J. 2000, *AJ*, 119, 136.
- Rosenberg, J. L. & Schneider, S. E. 2003, *ApJ*, 585, 256.
- Rosenberg, J. L. & Schneider, S. E. 2002, *ApJ*, 567, 247.
- Rosenberg, J. L. & Schneider, S. E. 2000, *ApJS*, 130, 177 [RS00].
- Scodreggio, M. & Gavazzi, G. 1993, *ApJ*, 409, 110.
- Scodreggio, M., Solanes, J. M., Giovanelli, R., & Haynes, M. P. 1995, *ApJ*, 444, 41.
- Spitzak, J. G. & Schneider, S. E. 1998, *ApJS*, 119, 159 [SS98].
- Shen, S., Mo, H. J., & Shu, C. 2002, *MNRAS*, 331, 259.
- Sprayberry, D., Impey, C. D., Bothun, G. D., & Irwin, M. J. 1995, *AJ*, 109, 558.
- Wevers, B. M. H. R., van der Kruit, P. C., & Allen, R. J., 1986, *A&SS*, 66, 505.
- Zavala, J., Avila-Reese, V., Hernández-Toledo, H., Firmani, C. 2003, *A&A*, 412, 633.
- Zwaan, M., Briggs, F. H., Sprayberry, D. & Sorar, E. 1997, *ApJ*, 490, 173.

Table 1. 2MASS Data for ADBS Galaxies

ADBS Num.	ADBS Name	α_{ADBS} deg	δ_{ADBS} deg	α_{2MASS} deg	δ_{2MASS} deg	R _{J21e} "	b/a _{J21}	J _{21e}	err _{J21e}	R _{Aper} "	J _{Aper}	err _{JAp}	Method.
rs001	000330+2312	0.8957	23.2011	0.895847	23.200817	20.2	0.58	12.189	0.031	50.0	11.644	0.050	V3
rs002	000407+2234	1.0292	22.5820	...	...	...	...	...	...	30.0	13.436	0.111	MI
rs003	000623+2347	1.5949	23.7891	1.595224	23.789404	14.2	0.36	14.347	0.085	25.0	13.661	0.125	V3
rs004	000900+2348	2.2269	23.8176	2.227971	23.816923	19.4	0.50	12.969	0.050	25.0	12.729	0.055	V3
rs005	001622+2237	4.0929	22.6170	4.093417	22.616209	13.4	0.42	13.609	0.046	25.0	13.371	0.089	V3
rs006	002249+2310	5.6894	23.1953	5.689276	23.195107	8.0	0.82	14.265	0.067	25.0	13.492	0.107	V3
rs007	002526+2136	6.3600	21.6040	6.361917	21.603350	...	...	...	...	15.0	14.9	...	MI
rs008	003426+2436	8.6108	24.6010	8.603510	24.603650	2.9	1.0	17.1	...	15.0	15.1	...	MI
rs009	003751+0838	9.4905	8.6348	9.490385	8.635223	36.7	0.7	10.821	0.025	70.0	10.432	0.029	V3
rs010	003811+2523	9.5458	25.3960	9.548830	25.395960	8.1	0.5	15.5	...	20.0	14.7	...	MI
rs011	004649+2134	11.7042	21.5830	...	...	2.8	1.0	17.1	...	7.0	17.0	...	MI
rs012	011440+2708	18.6679	27.1350	18.690270	27.135980	...	...	...	...	15.0	15.1	...	MI
rs013	014206+1235	25.5397	12.6008	25.540237	12.601927	32.4	0.56	10.952	0.021	60.0	10.740	0.032	V3
rs014	014246+1309	25.6950	13.1550	...	...	28.8	0.44	12.5610	0.036	70.0	11.954	0.075	V2
rs015	014527+2531	26.3650	25.5220	...	...	7.0	1.0	15.7930	0.228	60.0	12.936	0.140	V2
rs016	014729+2719	26.8757	27.3316	26.874540	27.333349	...	...	...	...	115.0 ¹	11.069	0.056	V3
rs017	014847+1034	27.1958	10.5690	27.218527	10.589932	30.2	0.34	11.793	0.025	30.0	11.690	0.025	V3
rs018	015011+2309	27.5492	23.1560	27.555040	23.159340	4.6	0.9	15.8	...	20.0	12.6	...	MI
rs019	015105+1235	27.7721	12.5880	...	...	...	...	...	...	30.0	13.869	0.130	MI
rs020	015434+2312	28.6429	23.2040	28.642710	23.204920	3.8	0.9	17.1	...	20.0	15.0	...	MI
rs021	015443+1033	28.6800	10.5660	28.692800	10.563700	...	...	...	...	...	...	...	NM
rs022	015906+2523	29.7888	25.3855	29.789383	25.385777	14.8	0.50	13.838	0.066	50.0	12.255	0.078	V3
rs023	020022+2434	30.0972	24.5805	30.097410	24.580576	24.7	0.76	11.816	0.029	60.0	11.461	0.050	V3
rs024	020045+2809	30.1883	28.1660	30.188300	28.166000	...	...	...	...	...	...	...	NM
rs025	020148+2632	30.4430	26.5466	30.443739	26.546976	20.7	0.48	12.013	0.023	30.0	11.895	0.029	V3
rs026	020320+1837	30.8342	18.6295	30.834126	18.629620	18.6	1.0	12.775	0.048	50.0	12.294	0.080	V3
rs027	020320+2345	30.8371	23.7610	30.837140	23.76093	5.8	1.0	15.9	...	30.0	14.2	...	MI
rs028	020405+2412	31.0208	24.2090	...	...	...	...	...	...	70.0	12.412	0.093	V2
rs029	020918+2534	32.3093	25.5706	32.309147	25.570778	20.8	0.52	11.803	0.021	40.0	11.521	0.031	V3
rs030	022807+1935	37.0462	19.5996	37.046249	19.599314	52.6	0.54	10.387	0.008	95.2 ¹	10.288	0.026	V2
rs031	022859+2808	37.2458	28.1450	...	...	...	...	...	...	30.0	14.352	0.223	MI
rs032	023322+2034	38.3442	20.5820	...	...	...	...	...	...	30.0	13.955	0.185	MI
rs033	023323+2810	38.3479	28.1790	38.317542	28.178069	...	...	...	...	...	...	...	NM
rs034	023432+2234	38.6358	22.5760	...	...	4.6	1.0	18.6	...	...	...	...	NM
rs035	023622+2526	39.0677	25.4238	39.067299	25.423820	38.8	0.4	11.651	0.017	135.1 ¹	10.844	0.033	V2
rs036	024328+2035	40.8671	20.5960	...	...	7.0	0.94	14.639	0.070	20.0	13.743	0.090	V2
rs037	025537+1938	43.9071	19.6340	...	...	2.0	1.0	17.6	...	...	...	...	NM
rs038	025726+1008	44.3608	10.1380	44.362360	10.431630	5.5	1.0	15.9	...	40.0	13.2	...	MI
rs039	025749+1110	44.4740	11.1831	44.473911	11.182944	25.5	0.70	12.134	0.041	60.0	11.772	0.073	V3
rs040	025942+2514	44.9245	25.2371	44.928493	25.241194	60.8	0.48	10.396	0.019	70.0	10.004	0.015	V3
rs041	030525+0835	46.3550	8.5870	...	...	7.0	0.52	16.043	0.190	30.0	14.170	0.201	V2
rs042	030546+2212	46.4450	22.2036	46.444927	22.203434	19.4	0.56	12.520	0.033	30.0	12.293	0.041	V3
rs043	031420+2409	48.5824	24.1539	48.582500	24.153975	9.5	0.68	14.462	0.073	25.0	13.869	0.132	V3
rs044	032327+1109	50.8505	11.1527	50.850002	11.152466	21.2	0.18	14.116	0.073	25.0	13.581	0.115	V3
rs045	033112+1410	52.7762	14.1770	52.776714	14.177110	18.8	0.44	13.081	0.046	30.0	12.849	0.071	V3
rs046	034444+0833	56.1920	8.5627	56.191986	8.562414	13.1	0.54	13.962	0.078	40.0	12.996	0.133	V3
rs047	033456+1507	53.7034	15.1421	53.703297	15.141906	12.9	0.30	14.726	0.084	20.0	14.358	0.168	V3
rs048	040325+2034	60.8550	20.5770	...	...	...	...	...	...	30.0	14.064	0.183	MI
rs049	040344+2209	60.9339	22.1582	60.933250	22.159042	29.2	0.80	10.836	0.020	60.0	10.673	0.029	V3
rs050	040411+2207	61.0405	22.1309	61.040314	22.131767	...	...	...	...	...	...	...	NM

Table 1—Continued

ADBS Num.	ADBS Name	α_{ADBS} deg	δ_{ADBS} deg	α_{MASS} deg	δ_{MASS} deg	R_{J21e} "	b/a _{J21}	J _{21e}	err _{J21e}	R_{Aper} "	J _{Aper}	err _{J_{Ap}}	Method.
rs051	040731+2210	61.8829	22.1690	61.885460	22.167440	11.8	1.0	12.0	...	15.0	11.7	...	MI
rs052	041255+1328	63.2051	13.4961	63.205006	13.495862	14.2	0.84	12.613	0.033	30.0	12.313	0.042	V3
rs053	041336+2528	63.4121	25.4829	63.411591	25.482941	49.7	0.28	10.815	0.016	50.0	10.682	0.017	V3
rs054	043622+1309	69.0929	13.1600	...	...	...	...	...	...	30.0	13.520	0.125	MI
rs055	043649+2409	69.2112	24.1522	69.210838	24.152411	12.8	0.38	14.393	0.086	25.0	13.698	0.138	V3
rs056	044310+1010	70.7942	10.1720	...	...	2.8	1.0	18.5	...	...	...	...	NM
rs057	044753+2346	71.9729	23.7800	...	...	3.0	1.0	18.4	...	...	...	...	NM
rs058	045254+2310	73.2258	23.1810	73.220940	23.166440	4.1	1.0	16.8	...	40.0	13.1	...	MI
rs059	051733+1934	79.4228	19.6000	79.422737	19.600281	30.1	0.22	12.553	0.033	30.0	12.470	0.048	V3
rs060	053017+2233	82.5729	22.5660	...	...	...	...	...	...	...	...	...	NM
rs061	053522+1332	83.8186	13.5489	83.818535	13.548637	19.9	0.32	12.950	0.030	30.0	12.783	0.054	V3
rs062	053844+1508	84.6671	15.1475	84.667244	15.147024	28.5	0.44	11.615	0.026	40.0	11.322	0.029	V3
rs063	055452+1509	88.6697	15.1712	88.669701	15.170404	24.9	0.46	12.295	0.034	60.0	11.793	0.070	V3
rs064	055517+2526	88.8221	25.4430	88.825625	25.442286	...	...	...	...	...	...	...	NM
rs065	060719+1936	91.8292	19.6030	91.831590	19.601240	8.5	0.7	14.550	...	25.0	13.7	...	MI
rs066	060733+0932	91.8892	9.5360	...	...	5.0	1.0	15.9	...	...	...	...	NM
rs067	061543+1110	93.9292	11.1670	93.937620	11.172750	10.0	0.4	14.3	...	25.0	13.3	...	MI
rs068	061729+2807	94.3552	28.1212	94.355087	28.121138	18.7	0.54	12.968	0.043	50.0	12.351	0.082	V3
rs069	062054+2008	95.2271	20.1410	95.263125	20.170111	...	...	...	...	15.0	15.4	...	MI
rs070	062103+2010	95.2629	20.1700	...	...	2.1	1.0	17.9	...	...	...	...	NM
rs071	062302+1108	95.7424	11.1417	95.742302	11.142012	45.1	0.36	12.414	0.087	40.0	12.216	0.078	V3
rs072	063549+1107	98.9550	11.1280	98.977410	11.136020	21.0	0.7	8.650	...	...	...	...	NM
rs073	063603+1109	99.0142	11.1520	98.975083	11.134497	...	...	...	...	...	...	...	NM
rs074	065008+2009	102.5622	20.1409	102.561920	20.140766	22.5	0.66	11.479	0.020	40.0	11.123	0.021	V3
rs075	065406+0834	103.5258	8.5720	103.523490	8.571820	6.7	0.4	16.1	...	20.0	14.8	...	MI
rs076	065718+1332	104.3250	13.5363	104.312889	13.539122	7.0	0.76	14.853	0.079	30.0	13.399	0.102	V2
rs077	070911+2036	107.3001	20.6023	107.299553	20.602835	20.9	0.76	11.350	0.019	40.0	11.131	0.023	V3
rs078	071028+2307	107.6204	23.1320	...	...	4.1	1.0	18.3	...	...	...	...	NM
rs079	071225+2342	108.1043	23.7124	108.104881	23.711788	...	...	...	...	30.0	12.935	0.083	NM
rs080	071352+1031	108.4671	10.5240	...	...	9.6	0.1	16.637	0.257	70.0	12.111	0.083	V2
rs081	071553+1207	108.9721	12.1170	108.969116	12.115514	7.0	1.0	15.159	0.209	60.0	12.749	0.134	V2
rs082	071831+2709	109.6334	27.1584	109.632622	27.157993	29.8	0.22	12.576	0.040	30.0	12.478	0.058	V3
rs083	072502+2348	111.2564	23.7832	111.256935	23.783270	32.3	0.54	11.088	0.019	50.0	11.002	0.021	V3
rs084	072507+0931	111.2931	9.5160	111.293152	9.515792	24.0	0.60	12.115	0.048	40.0	11.803	0.057	V3
rs085	072858+2035	112.2254	20.5911	112.225601	20.591793	32.1	0.40	11.712	0.027	50.0	11.472	0.040	V3
rs086	073445+2234	113.7050	22.5765	113.704926	22.576496	10.6	0.60	14.447	0.075	25.0	13.604	0.101	V3
rs087	073533+1131	113.8908	11.5190	...	...	7.0	1.0	15.651	0.219	50.0	13.096	0.147	V2
rs088	081538+2107	123.9100	21.1310	123.932500	21.131500	4.9	1.0	15.8	...	15.0	15.1	...	MI
rs089	081617+2308	124.0729	23.1391	...	...	7.0	1.0	15.582	0.154	50.0	14.400	0.369	V2
rs090	081707+2433	124.2808	24.5640	124.282700	24.562860	2.6	1.0	17.300	...	10.0	15.6	...	MI
rs091	081726+2110	124.3560	21.1637	124.355927	21.163906	11.9	0.54	14.545	0.087	30.0	13.757	0.148	V3
rs092	081821+2431	124.5822	24.5269	124.582169	24.526777	7.0	0.82	14.309	0.058	25.0	13.883	0.147	V3
rs093	081904+2111	124.7577	21.1859	124.757904	21.185919	21.8	0.84	12.259	0.037	60.0	11.805	0.065	V3
rs094	081915+2030	124.8034	20.5109	124.803284	20.510710	32.5	0.64	11.006	0.019	60.0	10.729	0.025	V3
rs095	081926+2345	124.8433	23.7472	124.843201	23.747368	10.3	0.60	14.149	0.062	25.0	13.757	0.124	V3
rs096	082430+1835	126.1290	18.5969	126.129059	18.597067	20.1	0.72	12.287	0.030	40.0	11.978	0.041	V3
rs097	082551+2807	126.4484	28.1182	126.448250	28.118372	22.0	0.26	13.699	0.067	25.0	13.405	0.096	V3
rs098	083209+2233	128.0474	22.5606	128.047058	22.560581	26.3	0.88	10.525	0.015	60.0	10.196	0.018	V3
rs099	084025+1834	130.1058	18.5670	130.113270	18.573810	4.6	0.4	17.200	...	30.0	14.1	...	MI
rs100	084316+1305	130.8162	13.0858	130.816666	13.085848	32.0	0.30	12.215	0.038	40.0	11.940	0.051	V3

Table 1—Continued

ADBS Num.	ADBS Name	α_{ADBS} deg	δ_{ADBS} deg	α_{2MASS} deg	δ_{2MASS} deg	R_{J21e} "	b/a _{J21}	J _{21e}	err _{J21e}	R_{Aper} "	J _{Aper}	err _{JAp}	Method.
rs101	084411+2208	131.0581	22.1369	131.057724	22.136482	7.5	0.74	14.726	0.100	25.0	14.089	0.221	V3
rs102	084427+0930	131.0918	9.5376	131.090469	9.537537	11.4	0.28	14.997	0.087	20.0	14.421	0.179	V3
rs103	084504+0932	131.2675	9.5400	131.281586	9.541821	...	...	...	...	15.0	15.552	0.344	V2
rs104	084743+1325	131.9239	13.4189	131.923630	13.419150	15.9	0.94	12.221	0.033	30.0	11.973	0.036	V3
rs105	084804+2519	132.0171	25.3320	132.014470	25.332790	3.8	0.6	17.3	...	7.0	16.4	...	MI
rs106	085751+1230	134.4658	12.5020	134.469040	12.495370	7.5	1.0	14.9	...	20.0	14.2	...	MI
rs107	085927+1108	134.8650	11.1390	134.847916	11.135059	9.6	0.72	14.661	0.091	25.0	13.982	0.146	V2
rs108	085953+2109	134.9744	21.1518	134.974915	21.151529	8.9	0.54	14.782	0.076	25.0	13.940	0.138	V3
rs109	090024+2536	135.1000	25.6130	135.098663	25.612085	18.8	0.86	14.247	0.076	60.0	12.740	0.106	V2
rs110	090102+1105	135.2631	11.0977	135.263443	11.098387	18.0	0.84	12.473	0.049	30.0	12.302	0.056	V3
rs111	090129+2110	135.3666	21.1755	135.365799	21.175022	9.9	0.74	14.305	0.071	25.0	13.744	0.118	V3
rs112	090143+1105	135.4207	11.0829	135.420670	11.082906	16.2	0.68	12.861	0.053	40.0	12.514	0.097	V3
rs113	090240+1306	135.6728	13.1074	135.672501	13.107930	16.0	0.38	13.544	0.049	30.0	13.351	0.108	V3
rs114	090526+2533	136.3597	25.5504	136.359344	25.551037	12.8	0.48	14.261	0.070	30.0	13.448	0.109	V3
rs115	090544+2532	136.4371	25.5390	136.437083	25.539000	...	...	...	...	...	...	...	NM
rs116	090548+2526	136.4496	25.4373	136.449554	25.437365	28.8	0.64	11.357	0.020	70.0	10.489	0.024	V3
rs117	090552+2520	136.4692	25.3440	136.469167	25.344000	...	...	...	...	7.0	16.6	...	MI
rs118	093314+2307	143.3181	23.1348	143.318192	23.134792	14.6	0.68	13.295	0.049	40.0	12.719	0.086	V3
rs119	093704+0932	144.2599	9.5403	144.259537	9.539845	13.7	0.32	14.009	0.073	25.0	13.454	0.128	V3
rs120	093805+0931	144.5335	9.5231	144.532791	9.523837	46.7	0.32	10.526	0.013	60.0	10.413	0.019	V3
rs121	094040+1205	145.1700	12.0970	...	...	...	...	...	...	30.0	14.101	0.203	MI
rs122	094259+0929	145.7213	9.4923	145.722443	9.494324	16.7	0.32	13.896	0.059	40.0	13.116	0.120	V3
rs123	094641+2407	146.6742	24.1240	...	...	7.0	1.0	15.423	0.171	50.0	13.729	0.255	V2
rs124	094744+2343	146.9544	23.7221	146.954193	23.722757	17.0	0.68	12.982	0.038	30.0	12.555	0.043	V3
rs125	095413+2806	148.5571	28.1000	...	...	1.8	1.0	18.200	...	...	...	...	NM
rs126	100352+1105	150.9700	11.0980	...	...	4.1	0.8	16.700	...	...	...	...	NM
rs127	100500+2132	151.2521	21.5380	...	...	10.4	0.34	15.427	0.117	40.0	14.728	0.363	V2
rs128	100508+2207	151.2871	22.1270	151.284830	22.119650	3.0	0.7	17.800	...	10.0	15.4	...	MI
rs129	100735+1306	151.8971	13.1100	151.888080	13.106690	6.0	1.0	16.000	...	20.0	14.9	...	MI
rs130	101421+2207	153.5906	22.1249	153.590363	22.124683	21.7	0.26	13.295	0.037	40.0	12.421	0.053	V3
rs131	101636+2107	154.1424	21.1229	154.142334	21.122868	29.3	0.80	10.293	0.009	40.0	10.217	0.007	V3
rs132	101903+2209	154.7650	22.1520	154.765000	22.152000	...	...	...	...	...	...	...	NM
rs133	102055+2009	155.2329	20.1540	155.233246	20.155014	7.0	0.26	16.929	0.297	60.0	14.033	0.346	V2
rs134	102517+2007	156.3242	20.1240	...	...	4.0	1.0	17.4	...	...	...	...	NM
rs135	102714+2531	156.8159	25.5187	156.815552	25.518274	7.0	1.00	15.377	0.113	25.0	14.424	0.182	V3
rs136	102830+1933	157.1250	19.5610	157.125830	19.561620	3.5	1.0	17.0	...	15.0	15.7	...	MI
rs137	102922+2605	157.3198	26.0993	157.320038	26.099096	49.5	0.26	10.891	0.013	30.0	10.993	0.010	V3
rs138	103143+2518	157.9292	25.3100	157.930374	25.308094	7.0	0.24	17.199	0.290	60.0	13.519	0.169	V2
rs139	103439+2305	158.6658	23.0900	158.666000	23.090010	1.1	1.0	19.3	...	40.0	14.0	...	MI
rs140	103542+2607	158.9250	26.1260	158.925232	26.126038	14.5	0.58	13.244	0.037	30.0	12.955	0.062	V3
rs141	103937+2519	159.9127	25.3225	159.912598	25.322615	15.6	0.82	12.380	0.034	40.0	12.008	0.053	V3
rs142	104208+2344	160.5459	23.7464	160.546112	23.746815	17.0	0.94	11.836	0.018	30.0	11.674	0.019	V3
rs143	104310+1330	160.7950	13.5130	160.773958	13.510869	...	...	...	...	30.0	14.8	...	MI
rs144	104702+2632	161.7614	26.5432	161.760864	26.542980	18.5	0.82	12.412	0.032	30.0	12.150	0.032	V3
rs145	104722+1404	161.8429	14.0700	161.845993	14.068090	7.0	1.0	15.822	0.164	70.0	12.153	0.053	V2
rs146	104829+1232	162.1221	12.5350	162.116226	12.533339	62.1	0.4	10.506	0.007	70.0	10.288	0.010	V2
rs147	104916+1226	162.3171	12.4350	162.322479	12.420360	12.0	0.36	15.359	0.109	50.0	13.963	0.209	V2
rs148	104954+1309	162.4677	13.1619	162.467484	13.161813	7.0	1.00	14.915	0.072	25.0	14.681	0.220	V3
rs149	105204+1008	163.0165	10.1482	163.015945	10.148239	33.8	0.78	11.041	0.032	70.0	10.288	0.030	V3
rs150	110331+1107	165.8829	11.1210	165.885040	11.120287	8.0	0.24	16.637	0.353	60.0	13.211	0.213	V2

Table 1—Continued

ADBS Num.	ADBS Name	α_{ADBS} deg	δ_{ADBS} deg	α_{2MASS} deg	δ_{2MASS} deg	R_{J21e} "	b/a $_{J21}$	J_{21e}	err $_{J21e}$	R_{Aper} "	J_{Aper}	err $_{J_{Ap}}$	Method.
rs151	110710+1834	166.7950	18.5720	166.791351	18.569765	...	...	...	...	50.0	13.451	0.161	V2
rs152	110742+1933	166.9250	19.5510	166.943451	19.549334	...	...	...	...	70.0	12.726	0.113	V2
rs153	111027+1007	167.6150	10.1260	167.605780	10.126110	5.0	1.0	16.3	...	15.0	15.0	...	MI
rs154	111032+1932	167.6342	19.5390	167.656250	19.538505	...	...	...	...	...	...	...	NM
rs155	111341+2131	168.4577	21.5213	168.457855	21.520327	10.7	0.52	14.643	0.088	25.0	13.454	0.095	V3
rs156	111852+1305	169.7330	13.0942	169.730515	13.092888	80.0	0.68	7.509	0.001	286.1 ¹	6.989	0.016	V2
rs157	112020+1259	170.0630	12.9932	170.062531	12.989819	80.0	0.74	7.516	0.002	279.6 ¹	6.835	0.016	V2
rs158	112026+1334	170.0706	13.5876	170.067947	13.590096	61.6	0.38	10.108	0.018	478.7 ¹	7.174	0.020	V2
rs159	112134+2010	170.4303	20.1691	170.429611	20.169542	79.1	0.64	9.544	0.018	105.0 ¹	9.431	0.016	V3
rs160	112914+2035	172.3138	20.5833	172.313721	20.583197	18.3	0.38	13.601	0.065	40.0	12.905	0.114	V3
rs161	112949+2207	172.4384	22.1268	172.438293	22.126610	18.8	0.76	12.588	0.041	40.0	12.280	0.060	V3
rs162	113115+2530	172.8418	25.5009	172.841888	25.501511	12.4	0.34	14.883	0.098	20.0	14.226	0.153	V3
rs163	113119+2306	172.8292	23.1160	172.838943	23.114342	7.8	0.18	16.774	0.237	50.0	14.331	0.378	V2
rs164	113816+1206	174.5540	12.1123	174.553879	12.112137	24.6	0.92	11.766	0.017	60.0	11.509	0.035	V2
rs165	113845+2008	174.6887	20.1430	...	...	10.2	0.4	14.2	...	...	...	...	NM
rs166	114101+1128	175.2441	11.4707	175.244736	11.471132	66.9	0.74	9.043	0.014	109.4 ¹	8.843	0.016	V3
rs167	114227+2007	175.6021	20.1192	175.602005	20.119471	17.4	0.66	12.123	0.024	30.0	11.918	0.029	V3
rs168	114921+2607	177.3279	26.1215	177.328110	26.121529	29.7	0.60	11.696	0.031	50.0	11.424	0.039	V3
rs169	115004+2628	177.5191	26.4800	177.518539	26.478996	36.4	0.48	11.035	0.015	60.0	10.837	0.022	V3
rs170	115040+2531	177.6679	25.5260	177.666140	25.526370	6.1	0.6	15.9	...	15.0	15.1	...	MI
rs171	115840+2519	179.6695	25.3158	179.669601	25.316420	34.8	0.22	11.879	0.020	30.0	11.809	0.023	V3
rs172	115843+2506	179.6817	25.1164	...	...	1.5	1.0	18.8	...	...	...	...	NM
rs173	115906+2428	179.7771	24.4690	179.774933	24.472343	7.0	1.0	16.276	0.285	40.0	14.739	0.394	V2
rs174	120033+2004	180.1362	20.0741	180.136734	20.073902	28.8	0.86	11.340	0.037	60.0	11.026	0.042	V3
rs175	120112+1406	180.2932	14.1045	180.293259	14.104513	27.8	0.28	12.552	0.030	40.0	12.279	0.051	V3
rs176	120351+2525	180.9740	25.4334	180.973862	25.434111	12.4	0.50	14.261	0.064	30.0	13.082	0.072	V3
rs177	121206+2518	183.0279	25.3120	183.027756	25.309816	7.0	1.0	16.379	0.330	25.0	15.390	0.474	V2
rs178	121233+1207	183.1347	12.1237	183.134674	12.123732	36.3	0.64	11.571	0.034	70.0	10.978	0.041	V3
rs179	121437+1205	183.6542	12.1000	183.637333	12.103053	...	...	...	...	...	...	...	NM
rs180	121514+1259	183.8199	13.0236	183.82016	13.023854	81.8	0.18	10.662	0.009	170.3 ¹	10.320	0.028	V2
rs181	121516+2038	183.8093	20.6593	183.809830	20.659077	35.7	0.28	12.605	0.050	60.0	11.593	0.057	V3
rs182	121555+1308	183.9745	13.1490	183.975906	13.149279	90.8	0.4	8.350	0.001	317.8 ¹	7.525	0.016	V2
rs183	122126+1130	185.3244	11.5107	185.324112	11.511108	50.5	0.40	10.947	0.033	101.2 ¹	10.580	0.044	V3
rs184	122140+1130	185.4199	11.5028	185.419010	11.502320	31.1	1.0	11.4	...	60.0	11.1	...	V3
rs185	122647+1133	186.6958	11.5550	186.714035	11.554358	7.0	1.0	15.681	0.291	30.0	15.237	0.828	V2
rs186	123119+1129	187.8456	11.4927	187.846405	11.492820	13.0	0.44	14.074	0.094	30.0	13.713	0.224	V3
rs187	123352+1510	188.4700	15.1760	188.453491	15.168147	7.9	0.8	13.602	0.026	70.0	11.543	0.038	V2
rs188	123546+2801	188.9891	27.9615	188.991333	27.959259	80.0	0.6	8.832	0.004	315.0 ¹	8.411	0.033	V2
rs189	123606+2602	189.0868	25.9873	189.086685	25.987391	...	...	...	...	325.6 ¹	7.161	0.016	V3
rs190	124020+1031	190.0694	10.5185	190.069168	10.518446	13.8	0.94	12.935	0.061	40.0	12.630	0.116	V3
rs191	124358+1307	190.9873	13.1268	190.98616	13.126577	93.0	0.58	8.949	0.005	190.5 ¹	8.647	0.020	V2
rs192	124412+1207	191.0521	12.1170	...	...	2.8	1.0	17.1	...	...	...	...	NM
rs193	124516+2708	191.3209	27.1252	191.321457	27.125490	26.9	0.70	11.409	0.021	50.0	11.139	0.027	V3
rs194	124820+0829	192.0962	8.4877	192.095566	8.487244	94.6	0.56	8.540	0.013	117.7 ¹	8.403	0.012	V3
rs195	124930+2528	192.3922	25.4696	192.392548	25.469921	37.9	0.44	11.475	0.028	70.0	11.022	0.044	V3
rs196	125029+2530	192.6100	25.5004	192.610886	25.500759	80.0	0.84	7.928	0.002	364.5 ¹	7.160	0.024	V2
rs197	125145+2607	192.9408	26.1180	192.935074	26.109131	7.0	1.0	16.422	0.444	50.0	13.696	0.257	V2
rs198	125156+1205	192.9799	12.0837	192.980713	12.083027	53.6	0.24	10.632	0.021	70.0	10.547	0.040	V3
rs199	125223+2138	193.0992	21.6370	...	...	1.8	1.0	18.3	...	...	...	...	NM
rs200	125411+2709	193.5199	27.1476	193.521866	27.149630	7.0	1.0	15.903	0.254	60.0	13.158	0.173	V2

Table 1—Continued

ADBS Num.	ADBS Name	α_{ADBS} deg	δ_{ADBS} deg	α_{2MASS} deg	δ_{2MASS} deg	R_{J21e} "	b/a $_{J21}$	J_{21e}	err $_{J21e}$	R_{Aper} "	J_{Aper}	err $_{J_{Ap}}$	Method.
rs201	125711+2135	194.1829	21.6821	194.183609	21.68528	90.8	1.0	6.886	0.001	308.8 ¹	6.267	0.016	V2
rs202	125850+1308	194.7108	13.1400	194.720306	13.152530	...	...	...	...	40.0	13.802	0.134	V2
rs203	130030+1007	195.1386	10.1301	195.138489	10.130094	11.6	0.92	12.880	0.042	30.0	12.688	0.073	V3
rs204	130543+2606	196.4131	26.1066	196.413406	26.106325	8.7	0.72	14.694	0.091	30.0	14.130	0.233	V3
rs205	130619+1025	196.5800	10.4320	196.593680	10.433570	...	...	...	...	40.0	13.4	...	MI
rs206	131051+1128	197.7356	11.4775	197.735382	11.477436	22.2	0.62	12.671	0.057	50.0	11.971	0.075	V3
rs207	131652+1232	199.2180	12.5487	199.218079	12.548311	15.7	0.90	12.723	0.044	40.0	12.315	0.067	V3
rs208	133643+0832	204.1829	8.5490	204.182570	8.548040	...	...	...	...	25.0	14.0	...	MI
rs209	135822+2533	209.5992	25.5499	209.599167	25.550037	15.4	0.34	14.029	0.069	25.0	13.632	0.120	V3
rs210	140122+1007	210.3252	10.1190	210.325211	10.118935	11.9	0.86	13.258	0.056	30.0	12.878	0.089	V3
rs211	141342+1230	213.4246	12.5041	213.424454	12.504229	17.7	0.72	12.470	0.040	40.0	12.171	0.064	V3
rs212	141449+2519	213.7166	25.3177	213.717957	25.317614	61.3	0.22	10.936	0.020	116.9 ¹	10.684	0.025	V3
rs213	141453+1407	213.7171	14.1254	213.716660	14.125856	24.8	0.48	11.878	0.024	40.0	11.620	0.032	V3
rs214	141556+2303	213.9871	23.0540	213.982178	23.053068	7.0	1.0	16.824	0.528	50.0	13.316	0.146	V2
rs215	142335+2131	215.8992	21.5250	...	...	2.6	1.0	16.8	...	...	...	...	NM
rs216	143307+1030	218.2882	10.5104	218.288239	10.510788	17.4	0.86	11.091	0.016	30.0	10.991	0.017	V3
rs217	143523+0930	218.8479	9.5020	218.845370	9.500750	5.5	0.9	16.0	...	10.0	15.4	...	MI
rs218	144842+1226	222.1774	12.4573	222.177338	12.457081	14.4	0.86	11.954	0.019	30.0	11.666	0.022	V3
rs219	145050+2519	222.7108	25.3280	222.710000	25.323839	...	...	...	...	...	...	...	NM
rs220	145647+0930	224.1971	9.5040	224.198715	9.509689	7.0	1.0	16.097	0.368	50.0	13.327	0.204	V2
rs221	145931+2706	224.8930	27.1164	224.893188	27.116291	24.1	0.36	12.687	0.044	50.0	12.067	0.077	V3
rs222	151612+1030	229.0457	10.5098	229.045197	10.509682	53.5	0.26	10.719	0.025	60.0	10.483	0.031	V3
rs223	152144+2309	230.4403	23.1525	230.440384	23.152630	17.8	0.76	13.017	0.050	40.0	12.386	0.062	V3
rs224	153438+1510	233.6621	15.1710	...	...	2.0	1.0	18.1	...	...	...	...	NM
rs225	153518+1203	233.8470	12.0473	233.846832	12.047511	47.6	0.66	10.484	0.027	70.0	10.286	0.029	V3
rs226	153703+2009	234.2619	20.1288	234.261551	20.128714	19.5	0.30	12.550	0.031	25.0	12.420	0.046	V3
rs227	154540+2805	236.4138	28.0887	236.414215	28.088715	28.6	0.30	11.809	0.023	30.0	11.733	0.028	V3
rs228	154540+1230	236.4200	12.5070	236.433456	12.510056	7.0	1.0	15.752	0.183	60.0	13.412	0.180	V2
rs229	154546+2033	236.4355	20.5602	236.435379	20.560698	16.5	0.78	12.998	0.054	50.0	12.102	0.075	V3
rs230	155946+1831	239.9421	18.5320	239.966830	18.529676	19.0	1.0	13.0	...	30.0	11.5	...	MI
rs231	160510+2032	241.3036	20.5419	241.303665	20.542391	7.4	0.68	11.684	0.021	40.0	11.546	0.030	V3
rs232	160556+0830	241.4850	8.5030	241.508830	8.506080	21.0	0.8	15.4	...	20.0	14.1	...	MI
rs233	161413+0932	243.5663	9.5376	243.566452	9.537125	15.9	0.88	12.031	0.028	30.0	11.901	0.034	V3
rs234	161830+2210	244.6248	22.1633	244.624588	22.163591	25.6	0.22	13.113	0.043	30.0	12.819	0.066	V3
rs235	163122+2010	247.8386	20.1850	247.836716	20.185495	42.8	0.20	11.859	0.029	40.0	11.727	0.038	V3
rs236	163130+2005	247.8750	20.0930	...	...	2.6	0.9	17.3	...	...	...	...	MI
rs237	163425+2132	248.6060	21.5408	248.606186	21.540836	40.2	0.38	10.722	0.015	60.0	10.496	0.021	V3
rs238	163455+2034	248.7306	20.5785	248.730484	20.578688	21.5	0.24	13.659	0.060	25.0	13.399	0.095	V3
rs239	164802+2133	252.0092	21.5610	252.009250	21.558161	...	...	...	...	10.0	16.2	...	MI
rs240	171634+2135	259.1417	21.5860	259.151860	21.585780	5.0	0.5	16.2	...	20.0	15.4	...	MI
rs241	172709+1132	261.7993	11.5447	261.798096	11.543621	53.6	0.32	10.294	0.017	60.0	10.228	0.020	V3
rs242	173438+2533	263.6582	25.5683	263.658142	25.568470	21.7	0.36	13.286	0.059	30.0	12.837	0.074	V3
rs243	173547+2533	263.9458	25.5540	263.957190	25.532180	2.6	0.6	16.1	...	25.0	14.4	...	MI
rs244	174924+1834	267.3754	18.5651	267.375671	18.565491	11.0	0.70	13.212	0.050	25.0	13.082	0.095	V3
rs245	175158+2134	268.0244	21.5692	268.024414	21.569176	27.7	0.60	11.766	0.030	50.0	11.391	0.038	V3
rs246	175845+2533	269.6812	25.5518	269.681274	25.551987	12.1	0.46	13.726	0.042	20.0	13.323	0.059	V3
rs247	180506+2308	271.2758	23.1470	...	...	2.2	0.7	18.0	...	...	...	...	NM
rs248	181221+1836	273.1344	18.5993	273.134277	18.599407	17.2	0.40	13.475	0.058	30.0	12.773	0.071	V3
rs249	181856+2708	274.7350	27.1430	...	...	6.0	1.0	16.3	...	...	...	...	NM
rs250	183229+2308	278.1225	23.1390	...	...	...	...	...	...	...	...	...	NM

Table 1—Continued

ADBS Num.	ADBS Name	α_{ADBS} deg	δ_{ADBS} deg	α_{2MASS} deg	δ_{2MASS} deg	R_{J21e} "	b/a _{J21}	J _{21e}	err _{J21e}	R_{Aper} "	J _{Aper}	err _{JAp}	Method.
rs251	184335+2011	280.8967	20.1913	...	...	5.0	1.0	16.3	...	...	...	...	NM
rs252	184829+1835	282.1221	18.5960	282.122480	18.597260	...	...	...	...	...	...	...	NM
rs253	192728+2012	291.8679	20.2060	291.867917	20.206000	...	...	...	...	...	...	...	NM
rs254	202344+0832	305.9371	8.5390	305.960780	8.538410	3.5	1.0	16.3	...	25.0	14.3	...	MI
rs255	210538+2709	316.4092	27.1520	316.406042	27.152461	...	...	...	...	7.0	16.4	...	MI
rs256	211113+2809	317.8050	28.1590	...	...	2.4	1.0	17.4	...	...	...	...	NM
rs257	214731+2209	326.8808	22.1650	326.879050	22.164010	3.6	0.6	17.1	...	10.0	15.7	...	MI
rs258	214813+2209	327.0547	22.1591	327.054321	22.159580	32.2	0.88	10.660	0.018	60.0	10.437	0.020	V3
rs259	223744+2347	339.4458	23.7863	339.445374	23.786671	74.7	0.22	9.745	0.012	107.6 ¹	9.693	0.012	V3
rs260	225557+2610	343.9879	26.1710	...	...	2.3	1.0	17.9	...	...	...	...	NM
rs261	230433+2709	346.1400	27.1560	346.140320	27.155710	14.4	0.44	14.643	0.104	25.0	14.008	0.151	V2
rs262	231941+1011	349.9228	10.1852	349.922363	10.184981	17.1	0.66	12.794	0.052	60.0	11.481	0.067	V3
rs263	234042+2613	355.1758	26.2320	355.166138	26.236441	8.2	0.36	15.789	0.172	70.0	12.106	0.081	V2
rs264	234734+1836	356.8950	18.6000	356.922119	18.600136	8.2	0.34	16.266	0.240	30.0	13.898	0.166	V2
rs265	235121+2034	357.8278	20.5778	357.827637	20.577906	13.6	0.40	14.137	0.064	25.0	13.729	0.117	V3

¹Sources significantly larger than 70'' have J-band extrapolated magnitudes and sizes substituted for the aperture values because the 70'' aperture values are severe underestimates of the sizes and fluxes of the systems.

Table 2. 2MASS Data for AS Galaxies

AS Name	α_{AS} deg	δ_{AS} deg	α_{2MASS} deg	δ_{2MASS} deg	R_{J21e} "	b/a_{J21}	J_{21e}	err_{J21e}	R_{Aper} "	J_{Aper}	err_{JAp}	Meas.
SS01	331.1152	23.3133	331.1197	23.3204	...	...	...	...	15.0	14.6	...	MI
SS02	337.0285	23.3737	...	...	...	...	...	...	7.0	17.7	...	MI
SS03	340.3504	23.3751	340.3521	23.3755	13.6	0.76	13.19	0.053	25.0	12.873	0.066	V3
SS04	344.7468	24.0036	344.7480	24.0038	15.3	0.26	14.28	0.079	25.0	13.837	0.159	V3
SS05	345.9213	23.6856	...	...	2.6	1.0	17.5	...	20.0	14.7	...	MI
SS06	347.0237	23.2605	...	...	...	...	...	...	...	...	...	NM
SS07	347.1364	23.6312	...	...	7.2	1.0	15.3	...	40.0	14.3	...	MI
SS08	350.2188	23.8071	350.2199	23.8074	33.9	0.16	12.04	0.021	25.0	12.040	0.026	V3
SS09	350.6530	23.3747	350.6534	23.3747	8.8	0.86	14.40	0.075	25.0	13.817	0.128	V3
SS10	351.0689	23.8434	...	...	...	...	...	...	...	...	...	NM
SS11	351.1405	23.9862	351.1415	23.9866	15.5	0.24	14.45	0.091	25.0	13.819	0.166	V3
SS12	351.9209	23.5888	351.9210	23.5889	24.7	0.62	11.66	0.027	40.0	11.411	0.031	V3
SS13	352.0248	23.5313	352.0260	23.5314	17.6	0.96	11.43	0.021	30.0	11.311	0.021	V3
SS14	352.9108	24.0364	...	...	...	...	...	...	...	...	...	NM
SS15	353.0082	23.5579	353.0090	23.5574	13.2	0.72	13.00	0.041	30.0	12.782	0.071	V3
SS16	353.1212	23.9311	353.1211	23.9310	28.5	0.68	10.80	0.016	50.0	10.690	0.021	V3
SS17	353.6124	23.9037	353.6132	23.9032	7.0	1.0	15.36	0.145	15.0	14.845	0.192	V2
SS18	353.6850	23.9713	...	...	...	...	...	...	...	...	...	NM
SS19	353.9638	23.6187	353.9651	23.6187	22.4	0.80	11.56	0.026	40.0	11.431	0.031	V3
SS20	355.8094	23.5113	355.8094	23.5118	15.9	0.30	13.85	0.054	25.0	13.692	0.109	V3
SS21	355.8252	23.5131	355.8263	23.5114	...	...	...	...	...	...	...	NM
SS22	0.4081	23.4845	0.4096	23.4836	41.9	0.38	10.62	0.017	70.0	10.100	0.022	V3
SS23	0.4232	23.4970	0.4247	23.4958	30.2	0.38	11.26	0.020	40.0	11.118	0.024	V3
SS24	1.2792	23.9720	1.2701	23.9694	7.0	0.6	15.3	...	20.0	14.7	...	MI
SS25	1.5939	23.7889	1.5952	23.7894	14.2	0.36	14.35	0.085	25.0	13.661	0.125	V3
SS26	2.2257	23.8174	2.2279	23.8169	19.4	0.50	12.97	0.050	25.0	12.729	0.055	V3
SS27	3.4943	23.9691	3.4941	23.9700	11.8	0.50	14.19	0.075	30.0	13.366	0.122	V3
SS28	4.7505	23.4782	4.7523	23.4765	12.2	0.64	13.66	0.056	30.0	13.115	0.092	V3
SS29	5.5197	23.7365	5.5212	23.7369	16.6	0.52	13.36	0.058	30.0	12.893	0.079	V3
SS30	6.4859	23.9238	...	...	...	...	...	...	...	...	...	NM
SS31	6.8649	23.7228	6.8655	23.7234	10.4	0.34	15.312	0.149	15.0	14.919	0.256	V2
SS32	8.1336	23.3954	8.1351	23.3951	14.1	0.42	14.04	0.077	30.0	13.181	0.108	V3
SS33	8.3984	23.3981	8.3996	23.3977	17.5	0.48	13.27	0.058	40.0	12.815	0.113	V3
SS34	8.9901	24.0374	8.9913	24.0370	13.5	0.36	14.35	0.083	25.0	13.740	0.141	V3
SS35	9.0151	23.9582	9.0168	23.9578	31.6	0.84	10.71	0.019	70.0	10.322	0.025	V3
SS36	9.2147	23.9914	9.2150	23.9909	32.6	0.76	10.39	0.015	60.0	10.196	0.018	V3
SS37	10.7042	23.4903	10.7041	23.4901	20.9	0.38	12.31	0.030	25.0	12.168	0.035	V3
SS38	10.7271	23.5031	10.7271	23.5030	20.8	0.66	11.83	0.027	40.0	11.503	0.034	V3
SS39	11.2149	23.8069	...	...	...	...	...	...	...	...	...	NM
SS40	12.4269	24.0459	12.4288	24.0455	9.2	0.74	13.47	0.043	20.0	13.154	0.065	V3
SS41	12.98	23.48	...	...	5.5	1.0	15.3	...	20.0	14.6	...	MI
SS42	14.0674	24.0538	...	...	...	...	...	...	...	...	...	NM
SS43	14.3303	23.8899	14.3321	23.8899	13.9	0.52	13.78	0.059	30.0	13.181	0.091	V3
SS44	19.9940	23.9652	19.9950	23.9647	7.0	1.0	15.216	0.130	20.0	14.449	0.182	V2
SS45	22.8405	23.9544	...	...	...	...	...	...	25.0	15.1	...	MI
SS46	23.7908	24.0077	...	...	6.6	1.0	15.2	...	15.0	14.9	...	MI
SS47	24.1481	23.8156	...	...	...	...	...	...	...	...	...	NM
SS48	26.2821	23.7699	...	...	...	...	...	...	15.0	15.8	...	MI
SS49	27.6763	23.4003	27.6777	23.3993	9.8	0.42	14.49	0.060	20.0	14.219	0.140	V3
SS50	28.4721	23.9116	28.4742	23.9141	...	...	...	...	15.0	14.7	...	MI
SS51	29.9746	23.6448	29.9769	23.6444	36.7	0.68	10.82	0.021	60.0	10.546	0.023	V3
SS52	30.1886	23.7543	...	...	...	...	...	...	10.0	15.9	...	MI
SS53	30.2624	23.4191	30.2634	23.4177	12.6	0.72	13.26	0.044	30.0	13.029	0.083	V3
SS54	30.8369	23.7580	30.8365	23.7612	3.9	1.0	16.8	...	50.0	13.6	...	MI
SS55	30.9358	24.0252	30.9296	24.0286	6.1	0.5	15.3	...	30.0	13.8	...	MI
SS56	32.2781	23.8435	...	...	...	...	...	...	20.0	15.5	...	MI
SS57	32.2964	23.2527	32.2977	23.2521	11.0	0.44	14.33	0.065	20.0	13.917	0.112	V3
SS58	33.4041	23.2470	33.4003	23.2585	2.5	0.6	17.1	...	10.0	15.8	...	MI
SS59	36.2982	23.8443	36.2925	23.8517	9.3	1.0	14.3	...	30.0	13.7	...	MI
SS60	36.3403	23.8285	36.3414	23.8272	11.7	0.80	13.33	0.047	30.0	13.026	0.084	V3
SS61	37.0907	23.2155	37.0923	23.2146	29.1	0.28	12.23	0.029	30.0	12.089	0.035	V3
SS62	37.1645	23.7972	37.1663	23.7964	15.0	0.66	12.65	0.032	25.0	12.355	0.036	V3
SS63	37.3371	23.9717	37.3392	23.9705	9.0	0.62	14.53	0.076	25.0	13.560	0.107	V3
SS64	38.1064	23.3647	38.1076	23.6525	3.1	0.8	16.6	...	15.0	15.2	...	MI
SS65	38.2242	23.3283	38.2242	23.3276	15.9	0.36	13.86	0.068	30.0	13.495	0.139	V3
SS66	38.3713	23.4862	38.3712	23.4850	12.6	0.36	14.783	0.098	25.0	13.887	0.147	V2

Table 2—Continued

AS Name	α_{AS} deg	δ_{AS} deg	α_{2MASS} deg	δ_{2MASS} deg	R_{J21e} "	b/a_{J21}	J_{21e}	err_{J21e}	R_{Aper} "	J_{Aper}	err_{JAp}	Meas.
SS67	38.6782	23.4142	38.6797	23.4131	38.6	0.74	10.43	0.017	60.0	10.251	0.017	V3
SS68	39.0389	23.9003	39.0404	23.8991	18.4	0.56	13.10	0.050	50.0	12.230	0.077	V3
SS69	39.3623	23.3013	39.3645	23.3000	15.6	0.62	12.86	0.042	40.0	12.468	0.079	V3
SS70	40.3270	23.2694	...	...	...	...	...	...	...	...	...	NM
SS71	41.7268	23.5962	41.7276	23.5956	20.2	0.58	12.00	0.023	30.0	11.858	0.026	V3
SS72	41.9639	23.4062	41.9655	23.4040	29.3	0.44	11.80	0.025	60.0	11.480	0.048	V3
SS73	42.1616	23.2747	...	...	...	...	...	...	...	...	...	NM
SS74	43.6756	23.3781	...	...	...	...	...	...	15.0	16.0	...	MI
SS75	44.3925	23.7871	...	...	...	...	...	...	20.0	15.9	...	MI

FIG. 1.— The ADBS galaxies in order of baryonic mass from the most massive to the least massive. (JPEG file included with download)

FIG. 2.— The ADBS galaxies for which baryonic mass was not able to be determined. (JPEG file included with download)

FIG. 3.— The AS galaxies in order of baryonic mass from the most massive to the least massive. (JPEG file included with download)

FIG. 4.— The AS galaxies for which baryonic mass was not able to be determined. (JPEG file included with download)

This figure "Rosenberg.figA1_comp.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0411499v1>

This figure "Rosenberg.figA2_comp.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0411499v1>

This figure "Rosenberg.figA3_comp.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0411499v1>

This figure "Rosenberg.figA4_comp.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0411499v1>

This figure "Rosenberg.figA5_comp.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0411499v1>

This figure "Rosenberg.figA6_comp.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0411499v1>